



Investigation of negative cloud radiative forcing over the Indian subcontinent and adjacent oceans during the summer monsoon season

B. V. Thampi, R. Roca

► To cite this version:

B. V. Thampi, R. Roca. Investigation of negative cloud radiative forcing over the Indian subcontinent and adjacent oceans during the summer monsoon season. *Atmospheric Chemistry and Physics*, 2013, 13 (11), pp.28895-28951. 10.5194/acpd-13-28895-2013 . hal-00996365

HAL Id: hal-00996365

<https://hal.science/hal-00996365>

Submitted on 26 May 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Negative cloud
forcing**

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Investigation of negative cloud radiative forcing over the Indian subcontinent and adjacent oceans during the summer monsoon season

B. V. Thampi^{1,*} and R. Roca²

¹Laboratoire de Météorologie Dynamique, UMR8539, UPMC, BP99, 4 place Jussieu, 75252 Paris, France

²Laboratoire d'Etudes en Géophysique et Océanographie Spatiales, UMR5566, OMP, 14 Avenue E. Belin, 31400 Toulouse, France

*now at: Science Systems and Applications, Inc. (SSAI), Hampton, VA, USA

Received: 15 August 2013 – Accepted: 12 September 2013 – Published: 6 November 2013

Correspondence to: B. V. Thampi (btlmd@lmd.jussieu.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Abstract

Radiative properties of clouds over the Indian subcontinent and nearby oceanic regions (0–25° N, 60–100° E) during the Asian summer monsoon season (June–September) are investigated using the Clouds and Earth's Radiant Energy System (CERES) Top of the Atmosphere (TOA) flux data. Using multi-year satellite data, the net cloud radiative forcing (NETCRF) at the TOA over the Indian region during the Asian monsoon season is examined. The seasonal mean NETCRF is found to be negative (with its magnitude exceeding $\sim 30 \text{ W m}^{-2}$) over (1) the northern Bay of Bengal (close to the Myanmar–Thailand coast), (2) the Western Ghats and (3) the coastal regions of Myanmar. Such strong negative NETCRF values observed over the Indian monsoon region contradicts the assumption that near cancellation between LWCRF and SWCRF is a generic property of all tropical convective regions. The seasonal mean cloud amount (high and upper middle) and corresponding cloud optical depth observed over the three regions show relatively large values compared to rest of the Indian monsoon region. Using satellite derived cloud data, a statistical cloud vertical model delineating the cloud cover and single scattering albedo was developed for the three negative NETCRF regions. The shortwave (SW), longwave (LW) and net cloud radiative forcing over the three negative NETCRF regions are calculated using the Rapid Radiative Transfer Model (RRTM) with cloud vertical model as input. The NETCRF estimated from CERES observations show good comparison with that computed using RRTM (within the uncertainty limit of CERES observations). Sensitivity tests are conducted using RRTM to identify the parameters that control the negative NETCRF observed over these regions during the summer monsoon season. Increase in atmospheric water vapor content during the summer monsoon season is found to influence the negative NETCRF values observed over the region.

ACPD

13, 28895–28951, 2013

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Radiation at the top of the atmosphere and surface of earth are significantly modulated by the presence of clouds in the atmosphere. Effect of clouds on the earth's radiation budget can be gauged by Cloud Radiative Forcing (CRF), which is defined as the difference between clear sky and total sky radiation (Charlock and Ramanathan, 1985). CRF can be estimated directly from satellite observation at the top of the atmosphere (TOA) (Ramanathan et al., 1989; Harrison et al., 1990) as well as at the surface by means of satellite inversion and modeling (Zhang et al., 1995; Rossow and Zhang, 1995; Pavlakis et al., 2008). Studies have shown that global average TOA shortwave cloud radiative forcing (SWCRF) is negative while TOA longwave cloud radiative forcing (LWCRF) is positive with net effect being cooling of the earth-atmosphere system. However, over tropical convective regions, TOA LW and SW cloud radiative forcing are found to cancel each other (Ramanathan et al., 1989; Kiehl and Ramanathan, 1990; Kiehl, 1994; Hartmann et al., 2001; Futyran et al., 2004). Kiehl (1994) theorized that occurrence of cloud tops close to tropical tropopause over the Western Pacific results in the observed near cancellation between the SW and LW CRF. Jensen et al. (1994) studied the sensitivity of TOA flux to cloud micro/macro-physical properties using a 1-D radiative transfer model and observed that net radiative forcing of cirrus near the tropical tropopause is positive for cloud optical depths less than ~ 16 and negative for larger optical depths. Studies show that changes in cloud particle size and optical depth can cause the NETCRF to vary between positive or negative values depending on the magnitude and direction of change (Zhang et al., 1999). Feedback of cloud radiative properties on the climate can be either positive or negative depending on the cloud microphysics and single-scattering properties (Liou and Ou, 1989; Stephens et al., 1990).

There exist few convective regions over tropics where near cancellation of SW and LW CRF is not strictly followed (Pai and Rajeevan, 1998; Rajeevan and Sreenivasan, 2000; Hartmann et al., 2001; Futyran et al., 2004; Balachandran and Rajeevan, 2007).

ACPD

13, 28895–28951, 2013

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Negative cloud forcing

B. V. Thampi and R. Roca

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Rajeevan and Sreenivasan (2000) has shown (using ERBE data) that net cloud radiative forcing is negative for a sizeable area over the Indian region during the summer monsoon season. Long-term studies show that the North Bay of Bengal region during the summer monsoon season is characterized by large amount of high clouds with cloud top altitude close to tropical tropopause (Devasthale and Fueglistaler, 2010; Meenu et al., 2010). The region also experiences highest integrated latent heat release observed over the planet during the Asian summer monsoon season (Zuluaga et al., 2010). Rajeevan and Srinivasan (2000) proposed that presence of large amount of optically thicker high level clouds are the main reason behind the observed negative NETCRF over the Bay of Bengal. Balachandran and Rajeevan (2007) showed that NETCRF over the oceanic regions of the Indian monsoon region are strongly correlated with changes in high cloud amount while over the land regions, both middle and high cloud amount variations make substantial contribution. Patil and Yadav (2005) observed that NETCRF over the Asian monsoon region undergo a year-to-year variability with maximum magnitude in 1988 and minimum in 1987 indicating association between monsoon rainfall activity and CRF over the region. Roca et al. (2004) studied the influence of atmospheric water loading on the LW cloud radiative forcing over the Bay of Bengal using idealized radiative transfer computations and showed that negative cloud radiative forcing is closely associated with increased atmospheric water vapor loading.

Earlier studies suggest that the negative NETCRF observed over the Indian region (0–25° N, 60–100° E) during the Asian summer monsoon season is mainly influenced by the cloud macro physical properties (cloud amount, cloud height) with little understanding on the impact of microphysical (cloud optical depth, particle size and habit) and environmental variables (water vapor, SST). A comprehensive study detailing the influence of these variables on NETCRF over the Indian region during summer monsoon season is yet to be carried out. Objective of the present study is to understand the influence of these parameters on the observed negative NETCRF over the Indian region during the summer monsoon season. This paper delineates various negative NETCRF regimes observed over the Indian region during the summer monsoon sea-

son using the Clouds and Earth's Radiant Energy System (CERES) TOA flux data and examines the factors that control it. Influence of cloud micro/macro physical properties and environmental variables on the cloud radiative forcing over the Indian region is analyzed using satellite observations and modeling. The paper is divided into following sections. Section 2 describes the data used in the estimation of CRF and error analysis. Section 3 examines the radiative transfer model used for the computation of TOA flux and different parameterization schemes. Section 4 examines the CERES observation of CRF over the Indian region. Section 5 describes the comparison of NETCRF from CERES and RRTM simulations and various sensitivity analysis performed. Sections 6 and 7 summarize the main results of the paper.

2 Data and methodology

2.1 CERES TOA flux data

The mean cloud radiative forcing over the Indian region during the summer monsoon season (June–September) of 2002–2005 was derived by analyzing the TOA flux data from CERES Aqua SRBAVG-GEO (Edition 2A) dataset, which contain the monthly mean regional TOA total-sky and clear-sky radiative fluxes (LW and SW) in a $1^\circ \times 1^\circ$ latitude/longitude grid. CERES SRBAVG-GEO data use narrow band radiance from geostationary meteorological satellites to account for changes in flux and cloud conditions between daily CERES observations, which reduce temporal sampling errors. The uncertainties in the estimated CERES TOA flux is relatively small compared to that derived from the ERBE data (Loeb et al., 2005) mainly due to better scene identification and incorporation of better angular distribution models (Smith et al., 2012). CERES SRBAVG clear-sky monthly mean TOA fluxes are derived from CERES Single Scanner Footprint (SSF) data that are completely cloud-free according to 1 km resolution Moderate Resolution Imaging Spectroradiometer (MODIS) data. However, due to the coarse spatial resolution of CERES instrument (20 km at nadir), estimation of

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



clear sky flux is possible only from geographical regions where clouds occur less frequently and cloud free regions have relatively large area. Because of this, clear-sky flux maps from CERES SRBAVG GEO contain many gaps (no clear sky flux data), especially over the Indian monsoon region. In order to circumvent this problem, TOA clear sky flux data from the CERES Terra Energy Balanced and Filled (EBAF) Edition 1A product (Loeb et al., 2009a) were used in the present study. CERES EBAF data provides TOA clear-sky fluxes for many overcast regions that have no CERES clear-sky observations over the course of a month. CERES SSF TOA clear-sky fluxes require 99 % of MODIS pixels (with 1 km nominal area) within the CERES footprint (20 km nominal) to be classified as clear. However, in a overcast cloudy region, there maybe 1 km clear-sky patch present inside a 20 km footprint region. The CERES EBAF product uses this clear-sky MODIS pixel radiance to derive broadband radiances, which are constrained to the overall CERES footprint to derive the TOA clear sky flux. In the present study, monthly mean TOA total sky flux from CERES SRBAVG GEO product and monthly mean TOA clear sky flux from the CERES EBAF product are used in the estimation of cloud radiative forcing over the Indian region during the summer monsoon season. More details regarding the CERES EBAF dataset are available online (http://ceres.larc.nasa.gov/documents/DQ_summaries/CERES_EBAF_Ed2.7_DQS.pdf).

CRF is used as a metric to assess the radiative impact of clouds on the climate system, which is defined as the difference between TOA clear and total-sky flux. Using the CERES TOA flux measurements, the cloud radiative forcing is calculated by taking the difference between TOA clear sky and total sky flux.

$$\text{SWCRF} = \text{SW}_{\text{clear}} - \text{SW}_{\text{total}} \quad (1)$$

$$\text{LWCRF} = \text{LW}_{\text{clear}} - \text{LW}_{\text{total}} \quad (2)$$

Where the subscripts “clear” and “total” represent the TOA clear-sky and total-sky fluxes, respectively. The net cloud radiative forcing (NETCRF) at the TOA is estimated by adding the shortwave and longwave cloud forcing.

$$\text{NETCRF} = \text{SWCRF} + \text{LWCRF} \quad (3)$$

Which can be re written as,

$$\text{NETCRF} = \text{SW}_{\text{clear}} + \text{LW}_{\text{clear}} - (\text{SW}_{\text{total}} + \text{LW}_{\text{total}}) \quad (4)$$

To determine the cloud radiative forcing over the Indian region, monthly mean TOA total-sky flux from CERES SRBAVG GEO and clear-sky flux from CERES EBAF dataset during the summer monsoon season (June–September) of 2002–2005 are used.

2.2 Uncertainty analysis

This section presents the methodology used to compute the total uncertainty in NETCRF values due to uncertainties in the CERES TOA flux measurement. Uncertainties in the CERES flux measurement can be broadly categorized into three main components; Sampling errors, calibration errors and algorithm errors. Sampling error refers to error associated with time sampling and spatial averaging of the data associated with the instrument normalization (Young et al., 1998), which corresponds to 0.3 W m^{-2} for the SW and LW flux (Loeb et al., 2009a). An additional error term equal to the standard error of the regional mean TOA flux is also added to the sampling error term. The term “algorithm error” refers to the errors associated with data retrieval which in this case mainly stems from uncertainties associated with narrow-to-broadband conversion of radiance, Angular Distribution Models (ADM) and scene identification whereas calibration error refers to the instrument measurement error (Level 1 product error). More details regarding the of CERES TOA flux estimation and associated uncertainties can be found in Loeb et al. (2009a) as well as in the CERES EBAF data quality summary document. In the present analysis, it is assumed that sampling, calibration and algorithm errors associated with the CERES TOA flux are uncorrelated and independent. The total uncertainty in the TOA flux due to these different error sources can be expressed as (Chambon et al., 2012),

$$\delta_{\text{TOAFLUX}} = \sqrt{\delta_{\text{sampling}}^2 + \delta_{\text{calibration}}^2 + \delta_{\text{algorithm}}^2} \quad (5)$$

28901

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Using the above equation, uncertainty in the CERES TOA clear and total sky flux (LW and SW) are estimated. The uncertainty computations are performed at the monthly mean scale using error estimates for a $1^\circ \times 1^\circ$ latitude/longitude grid area.

In order to estimate the uncertainty in NETCRF, effect of propagation of variable uncertainty on the uncertainty of a function is to be considered. From Eq. (4), it can be seen that NETCRF is a function of SW and LW flux. Hence total uncertainty associated with the CERES NETCRF measurement is related to uncertainties in CERES SW and LW flux (Loeb et al., 2009b). Using the general law of error propagation, it is possible to analytically determine how measurement uncertainty propagates into quantities, which are functions of the measurement. For a multi-variable function $y(x_1, x_2, x_3, \dots, x_N)$, the total uncertainty in y due to uncertainty in the input variables x (assuming that error contributions are small compared to the absolute value of the variable) can be expressed as (Taylor, 1982; Lo, 2005)

$$\delta y = \sqrt{\sum_{i=1}^N \left(\delta x \frac{\partial y}{\partial x_i} \right)^2 + \sum_{i \neq j=1}^N \left(R_{ij} \delta x_i \delta x_j \frac{\partial y}{\partial x_i} \frac{\partial y}{\partial x_j} \right)} \quad (6)$$

where δy is the total uncertainty in y , δx_i and δx_j are the uncertainties associated with the input variables (x_i and x_j) and R_{ij} represent the correlation coefficient between the input variables. The uncertainty in y are governed by the (a) change in y for a given change in the variables x_i and x_j (partial derivatives), (b) uncertainties in the input variable δx_i and δx_j and (c) how the variables x_i and x_j are correlated. If x_i and x_j are not correlated and independent of each other, second term in Eq. (6) vanishes and the equation takes the form of a Gaussian error propagation formula (Taylor, 1982; Evans et al., 1984). Depending on the correlation between individual input variables and sign of the product of partial derivatives in the second term of Eq. (6), uncertainty in y can increase or decrease. The uncertainties in the CERES TOA SW and LW (clear and total sky) flux represent the input variables in the Eq. (6) and R represent the correlation between these two variables. However, if TOA flux from two different dataset

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



were used for computing the NETCRF (eg., total-sky flux from CERES and clear-sky flux from ECMWF reanalysis), the input fluxes will not be correlated and the second term in Eq. (6) will vanish. In the present analysis, uncertainty in the CERES TOA SW and LW flux (using Eq. 5) as well as correlation coefficient between the fluxes are computed. Using these values as input in Eq. (6), the total uncertainty in the NETCRF is estimated.

2.3 Cloud data

The seasonal mean cloud parameters from the CERES SRBAVG2-GEO dataset for the summer monsoon season of 2002–2005 are used in the study. The SRBAVG2 GEO cloud data include layer averaged monthly mean cloud and aerosol retrievals from MODIS and geostationary satellites (Remer et al., 2005; Menzel et al., 2008). The MODIS derived cloud fraction and cloud optical depth are believed to be more accurate than that from geostationary satellites owing to the higher quality of the MODIS data. Studies show that MODIS could detect cirrus clouds over tropics having cloud optical depth as low as 0.02 with an uncertainty factor of 2 (Dessler and Yang, 2003). However, combining geostationary satellite data measurements with the CERES-MODIS data minimizes the temporal sampling errors and incorporates the diurnal variation in cloud amount and TOA flux between the two daily MODIS Aqua measurements. The SRBAVG2-GEO cloud data provides information about retrieved cloud parameters in four atmospheric layers that are combination of MODIS and geostationary derived cloud retrievals. Seasonal mean values of cloud fraction, cloud top height, cloud particle size and cloud optical depth available from SRBAVG GEO cloud database for high (cloud top pressure $h < 300$ hPa), upper middle ($h = 300$ – 500 hPa), lower middle ($h = 500$ – 700 hPa) and low level ($h > 700$ hPa) clouds respectively, are used in the analysis.

2.4 ISCCP FD TOA flux data

The International Satellite Cloud Climatology Project (ISCCP) data is an archive of more than 20 yr of global cloud observations. ISCCP data utilizes the radiance information from a series of geostationary satellites to create 3 hourly global maps of cloudiness. The ISCCP-FD data is an improved version of a previous ISCCP radiative flux product (Zhang et al., 1995) and provides radiative fluxes at the TOA, surface, and several levels within the atmosphere (Zhang et al., 2004). ISCCP FD data provides global total-sky and clear-sky fluxes (at surface, 680 mbar, 440 mbar, 100 mbar and TOA) for every 3 h interval in the shortwave and longwave range. Inter comparison of monthly mean fluxes from ISCCP-FD with ERBE and CERES suggest that there exist an uncertainty of the order of $\sim 5\text{--}10\text{ W m}^{-2}$ in the calculated TOA flux from ISCCP-FD dataset. Details regarding the data and methodology adopted in the estimation of ISCCP FD flux are provided in Zhang et al. (2004) and Rossow et al. (2005). In the present study, TOA SW and LW fluxes from the ISCCP radiative flux dataset (ISCCP-FD) for the June–September months of 2002–2005 are used to estimate the seasonal mean CRF over the Indian region.

3 Rapid Radiative Transfer Model (RRTM)

The rapid radiative transfer model is a band model for the calculation of longwave and shortwave atmospheric radiative fluxes and heating rates (Mlawer et al., 1997; Iacono et al., 2000; Clough et al., 2005). RRTM use the correlated- k method, which is a accurate and computationally fast radiative transfer scheme. TOA LW fluxes calculated by RRTM agree with those computed by line-by-line radiative transfer model within $\sim 1\text{ W m}^{-2}$ range, while SW fluxes agree within $\sim 1.5\text{ W m}^{-2}$ range (Clough et al., 2005; Morcrette et al., 2008). During the Spectral Radiance Experiment (SPECTRE) program, results from the RRTM flux simulations are validated against other radiation models for various atmospheric conditions (tropical, mid-latitude summer/winter) (Ellingson

ACPD

13, 28895–28951, 2013

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and Fouquart, 1991; Ellingson and Wiscombe, 1996). An important feature of RRTM is that it incorporates Monte-Carlo Independent Column Approximation (McICA) to represent sub-grid cloud variability (Barker et al., 2003; Pincus et al., 2003). A 1-D column version of RRTMG is used in the present study to compute the TOA fluxes. RRTM input data typically consists of surface emissivity, cloud/aerosol optical depth, altitude profile (60 atmospheric pressure levels) of temperature, pressure, cloud fraction and single scattering albedo, atmospheric mixing ratio profiles of water vapor, ozone, CO₂, methane and other trace gases, etc.

In the present study, a tropical model atmosphere incorporating above mentioned parameters are used as input in RRTM simulations. An idealized altitude profile of atmospheric water vapor mixing ratio is built and used in the model simulations. For this purpose, altitude profile of relative humidity (RH) in the atmosphere is constructed following the methodology adopted in Roca et al. (2004). In order to construct the vertical profile, first it is assumed that the relative humidity is constant in the atmospheric boundary layer (between surface and 850 hPa) and in the stratospheric layers (< 100 hPa). RH value of 85 % is assumed for the boundary layer and 10 % for the stratospheric layer in all the profiles. The free tropospheric (between 850 hPa and 100 hPa) humidity is then varied from 5 % to 100 % in steps of 5 to create a number of RH altitude profiles. By converting the relative humidity to corresponding water vapor mixing ratio, a number of idealized water vapor mixing ratio profiles are created. The precipitable water (PW) concentration for each profile is estimated by integrating the water vapor mixing ratio from surface to top of the model atmospheric layer. In the present model simulations, changing the free tropospheric humidity (from 5 % to 100 %) in the model atmospheric profile leads to a corresponding change in the PW from 33 to 68 mm.

Using RRTM, TOA flux is estimated for every half hour interval for full day, which is then integrated to obtain the daily mean TOA SW and LW flux. The total sky SW and LW flux are estimated by incorporating cloud parameters (cloud cover, optical depth and single scattering albedo) in to the model computation, whereas clear sky flux estima-

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tion do not include any cloud data. In the computation of total sky LW flux using RRTM, altitude profile of mean cloud cover and cloud optical depth (from CERES data) for the study region is used. For the total sky SW flux computation, in addition to cloud cover and optical depth, altitude profile of single scattering albedo is also required (which is not available from the satellite measurements). In addition, the model also requires information regarding vertical overlap between different cloud layers. However, satellite derived cloud cover data do not provide information about overlap among different cloud layers in the atmosphere without which modeling of TOA flux is difficult. To circumvent this deficiency, parameterization schemes are used to determine the vertical overlap between different cloud layers and cloud single scattering albedo (ice and water clouds). Details regarding the cloud overlap scheme and single scattering albedo parameterizations are described in the subsequent sections.

3.1 Parameterization schemes

3.1.1 Cloud overlap

Modeling of radiative flux due to clouds are complicated by difficulties in parameterizing its single-scattering properties (Liou, 1986) and cloud vertical structure (Weare, 1999; Rossow et al., 2005). Satellite observations of multi-layered clouds from space only provide information about the topmost cloud layer encountered with lower level clouds being either fully or partially observed. If there is no overlap among different cloud layers in an atmospheric column, then cloud amount at each level observed by the satellite is the actual cloud amount. However, this assumption does not hold true in most cases involving partially cloudy skies. When there is overlap among different cloud layers, information about actual amount of cloud in the lower levels are not fully recorded by satellites. Even with enhanced satellite and surface observation capabilities, information about cloud vertical structure is rather limited. Presence of partially filled cloud layers in the atmosphere creates problem in the model computation of the radiative fluxes because of the nonlinear relation between cloud properties and TOA

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



fluxes (Bergmann and Hendon, 1998). This, along with lack of information about the vertical cloud overlap in the model can lead to large errors in the estimated radiative fluxes. In order to circumvent this problem, most models employ cloud overlap schemes for computing the radiative fluxes (Collins, 2001; Zhang et al., 2004; Rossow et al., 2005; Cole et al., 2011). Cloud overlap schemes are also used by satellite simulators in comparing the simulated cloud data with observations from passive or active remote sensing instruments (Klein and Jakob, 1999; Webb et al., 2001; Zhang et al., 2005; Bodas-Salcedo et al., 2011). Most radiative transfer models use vertical cloud overlap schemes like maximum overlap, random overlap or combination of maximum and random overlap between cloud layers. In the present study, a cloud vertical model is developed using a type of maximum/random cloud overlap scheme.

Geleyn and Hollingsworth (1979) theorized that if clouds appear in two adjacent atmospheric layers, such cloud layers are usually vertical parts of the same cloud and there should be maximum overlap between them. Maximum overlap between two cloud layers can be expressed mathematically as,

$$C_{\max} = \text{Max.}(C_n, C_{n-1}) \quad (7)$$

Where C_n and C_{n-1} represents the cloud fraction of two adjacent cloud layers and C_{\max} represent total cloud amount due to the overlap of two cloud layers. The random overlap assumption holds good only when the cloud layers are separated by at least one clear-sky layer. The random cloud overlap scheme assumes that the cloudiness in any given cloud layer is independent of the cloudiness of other layer (Warren et al., 1985). The total cloud amount in a vertical column assuming random overlap between cloud layers can be expressed mathematically as (Stephens et al., 2004),

$$C_{\text{total}} = 1 - \prod_{n=1}^I (1 - C_n) \quad (8)$$

Where C_{total} is the total cloud amount, C_n is cloud fraction for a given cloud layer n . Tian and Curry (1989) showed that clouds tends to follow the maximum overlap scheme for

small horizontal cloud area whereas they follow the random overlap for large horizontal area ($\geq 500 \text{ km}^2$). But in reality, there are no completely random or maximum cloud overlap occurrences in nature but rather specific combinations of cloud types associated with specific meteorological conditions (Hahn et al., 2001; Rossow et al., 2005).

5 Whether observed from satellites or surface, there exists a specific overlap relationship among different cloud types for each meteorological situation. In the present study, we are trying to develop a cloud overlap scheme that represent the altitude structure of a convective cloud system characterized by contiguous cloud layers. Since contiguous cloud layers can be expected to possess fairly high degree of vertical correlation, a combination of random and maximum overlap schemes are used to represent the cloud vertical structure. Here it is assumed that cloud layers belonging to a particular cloud block (eg., all the cloud layers in the high cloud group) in a convective system are maximally overlapped, whereas adjacent cloud blocks are randomly overlapped (eg., between high and upper middle cloud group). Therefore, effective cloud fraction for all the cloud layers belonging to a particular cloud block will remain same (due to maximum overlap) whereas it will change from one cloud block to another (due to random overlap between two adjacent cloud blocks). Chou et al. (1998) also adopted a similar type of maximum/random assumption with maximum cloud overlap in each of three cloud regions (lower, middle, and upper troposphere) and random overlap between these cloud regions. Using this methodology (Eqs. 7 and 8), altitude profile of cloud cover is constructed using the cloud fraction data. A graphical representation of typical altitude structure of contiguous cloud layers calculated using the cloud overlap scheme is shown in Fig. 1.

25 The cloud vertical model is built as follows: the CERES GEO layer averaged cloud properties are defined for four cloud groups mainly; high, upper middle, lower middle and low level clouds. The mean cloud top pressure for each cloud group defines the boundary of cloud blocks in the model atmosphere; ie, cloud layers belonging to high cloud block in the model atmosphere are defined between the CERES mean high cloud top pressure (usually between 180–250 hPa in the model) and upper middle cloud top

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



pressure (between 300–400 hPa in the model). In the present cloud overlap scheme, the effective cloud fraction for all high cloud layers in the model will be equivalent to the CERES high cloud fraction since all cloud layers within a particular cloud block are maximally overlapped. Effective cloud fraction for the upper middle cloud block in the model is estimated assuming random overlap (using the Eq. 7) between the CERES high cloud and upper middle level cloud fraction. This newly estimated cloud fraction (using Eq. 8) is assigned to all the layers in the upper middle cloud block (maximum overlap) in the model atmosphere defined between the upper middle cloud top pressure and lower middle cloud top pressure. Using this methodology, effective cloud fraction for lower middle and low level clouds are also estimated. Base of the newly constructed cloud vertical profile is fixed at the top of boundary layer (850 hPa) while the cloud top coincides with that of the high level cloud.

3.1.2 Cloud single-scattering albedo (SSA)

For estimating TOA SW flux using RRTM, altitude profile of cloud SSA is required along with cloud cover information. Single scattering properties of clouds are governed by the cloud particle size, shape and water content. They also vary over a large range of values depending on the wavelength band under consideration. In this section, parameterization schemes used for deriving the single scattering albedo (SSA) of ice and water clouds are explained. SSA parameterization provides a mathematical relationship between the cloud properties (particle size, optical depth, water content) and single scattering albedo. Several attempts were made to parameterize the SSA of a cloud system solely based on the cloud water content alone (Sun and Shine, 1994; Platt, 1997). However, it was observed that cloud water content alone is insufficient and information about the cloud particle size is also required in the parameterization of SSA (Wyser and Yang, 1998). For water clouds, the single-scattering properties can be effectively parameterized either in terms of their average size (Slingo, 1989; Hu and Stamnes, 1993) or based on the water content/optical depth (Fouquart and Bonnel, 1980; Fouquart, 1985; Räisänen, 1999). Using the parameterization methodology of

Fouquart (1985), single scattering albedo (ω) for water clouds are estimated for different values of cloud optical depth. The parameterization equation can be expressed as (Fouquart, 1985),

$$\omega = 1 - (9 \times 10^{-4} + 2.75 \times 10^{-3}(\mu + 1) \exp(-0.09\tau)) \quad (9)$$

5 Where μ is the solar zenith angle and τ is the cloud optical depth. This equation is based on calculations for a lower-tropospheric cloud with a specific drop size distribution having effective radius of 9.9 μm . Using the above formula, SSA values were calculated for water clouds with varying optical depth and used in the present study. Unlike water clouds, spherical particle assumption is not valid in determining the single scattering properties of ice clouds since they take on a variety of shapes like plates, hexagonal crystals, bullet rosettes (Schmidt et al., 1995). Because of this, single-scattering
10 albedo of ice clouds are defined mainly by the effective cloud particle size (Hu and Stamnes, 1993; Fu, 1996). The effective particle size of a ice cloud can be defined mathematically as,

$$15 \quad R = 3/4 \frac{\int_{L_{\min}}^{L_{\max}} V(L)n(L)dL}{\int_{L_{\min}}^{L_{\max}} A(L)n(L)dL} \quad (10)$$

Where L is the dimension of an ice crystal, $V(L)$ is the volume of the crystal, $A(L)$ is the projected area, and $n(L)$ is the size distribution. The parameterization scheme takes into account the effective size of ice crystal, which removes the ambiguity regarding the particle shape and size from the SSA estimation. The single scattering albedo of
20 an ice crystal can be expressed as (Key et al., 2002),

$$\omega_i = b_0 + b_1 R + b_2 R^2 + b_3 R^3 \quad (11)$$

Where b_0 , b_1 , b_2 and b_3 are the empirical coefficients determined through regression for different SW spectral bands (i) and R is the mean ice particle size (estimated from

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



CERES SRBAVG2 GEO data). The bulk single scattering albedo (ω) of ice crystals for the entire SW band is computed by integrating the individual SSA values computed for different shortwave bands and expressed mathematically as (Slingo and Schrecker, 1982; Chou et al., 1998),

$$\omega = \frac{\sum \beta_i(\lambda) \omega_i(\lambda) S(\lambda)}{\sum \beta_i(\lambda) S(\lambda)} \quad (12)$$

Where β is the mean extinction coefficient and S is normalized irradiance in each spectral band. The above parameterization scheme for ice cloud single-scattering albedo is an extension of the Streamer radiative transfer model (Key and Schweiger, 1996), which has been validated for different ice crystal size distributions and habits. Using this method, SSA values are computed for ice cloud particles and used in the computation of TOA flux. Using these SSA parameterization schemes, a cloud vertical model delineating the effective cloud cover and SSA are developed and used to model the TOA flux and CRF.

4 Radiative characteristics of convective clouds over the Indian region

Figure 2a and b show the seasonal mean variation of SWCRF and LWCRF respectively, over the Indian region during the summer monsoon season of 2002–2005 derived from the CERES TOA flux data. From Fig. 2, it can be seen that large positive LWCRF values ($> 60 \text{ W m}^{-2}$) and negative SWCRF values ($< -100 \text{ W m}^{-2}$) are observed over oceanic (north Bay of Bengal and northeast Arabian Sea) and land regions alike (coastal Myanmar, Thailand, Cambodia, inland regions of China and Western Ghats). Absolute magnitude of SWCRF is greater than LWCRF over regions where LWCRF values exceeds 60 W m^{-2} (similar observations were also reported by Rajeevan and Sreenivasan (2000) using ERBE data). Absolute values of LWCRF and SWCRF show relatively lower values over rest of the oceanic and land areas. Figure 3 shows the seasonal mean variation of NETCRF over the Indian region during

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the summer monsoon season of 2002–2005. From Fig. 3, it can be seen that seasonal mean NETCRF show large variation over the Indian region with values ranging from $+30 \text{ Wm}^{-2}$ to -80 Wm^{-2} . No specific pattern exists in the regional variation of NETCRF over the Indian region. Based on the seasonal mean variation of NETCRF, the Indian region can be categorized into three distinct NETCRF regimes; (a) positive NETCRF regime ($> 20 \text{ Wm}^{-2}$) over the south Indian land mass as well as over the Srilankan region (b) near zero NETCRF (between $+20$ and -20 Wm^{-2}) over the oceanic regions to south of Indian land mass as well as over northern India and (c) negative NETCRF regime ($< -20 \text{ Wm}^{-2}$) over the north Bay of Bengal close to Myanmar coast, Bangladesh, Myanmar, Inland china and over the northeast Arabian Sea as well as over the Western Ghats. In the present study, we focus mainly on the negative NETCRF regimes over the Indian region (delineated by black boxes in Fig. 3). They are designated as, (1) the Bay of Bengal ($10\text{--}22^\circ \text{ N}$, $85\text{--}100^\circ \text{ E}$) region representing oceanic regime, (2) Myanmar ($15\text{--}20^\circ \text{ N}$, $92\text{--}100^\circ \text{ E}$) and (3) the Western Ghats ($10\text{--}20^\circ \text{ N}$, $72\text{--}77^\circ \text{ E}$) representing land regime.

Table 1 presents the seasonal mean TOA flux and CRF estimated from CERES data for the three regions during summer monsoon season of 2002–2005. In general, clear sky SW and LW flux values are relatively lower over the Bay of Bengal region. The mean SWCRF and LWCRF values are larger (in magnitude) over the Bay of Bengal (BOB) compared to Myanmar (MYN) and the Western Ghats (WGS). In all the three regions, the absolute value of SWCRF is larger than LWCRF indicating imbalance between the two. The mean NETCRF estimated for the three regions vary between -30 Wm^{-2} and -37 Wm^{-2} with largest values (in absolute magnitude) observed over Myanmar region. However, mean NETCRF values estimated for the three regions (shown in Table 1) do not bring out the complete picture without quantifying the uncertainties associated with it. Using methodology described in Sect. 2.2, the uncertainty associated with the CERES TOA flux and NETCRF for the three regions are computed and presented in Table 2 and 3, respectively. Table 2 presents various uncertainties associated with CERES TOA flux values over the three regions. Using these values as

input to Eq. (6), total uncertainty in the NETCRF for the three regions are computed and presented in Table 3. In general, the total uncertainty in the estimated NETCRF from CERES TOA flux measurements varies between 3 and 6 Wm⁻². Estimated uncertainty values are smallest for the Bay of Bengal region and largest for the Western Ghats. It must be noted that total uncertainty in the CERES NETCRF measurements are much lower compared to that from ERBE NETCRF measurements (Wielicki et al., 1995; Cess et al., 2001).

4.1 Cloud characteristics over the Indian region

Figure 4a and b depict the seasonal mean variation of cloud area fraction (in percentage) for the high and upper middle level clouds over the Indian region during the summer monsoon season estimated from CERES cloud data. Only high and upper middle level cloud fractions are shown in Fig. 4, since fractional coverage of lower middle and low level clouds are relatively lower (~ 10 % or less) over the most of the negative NETCRF regions. From Fig. 4a, it can be seen that high cloud fraction is relatively large (> 40 %) over the North Bay of Bengal and land areas over eastern India compared to that observed (~ 20–30 %) over oceanic regions to the south and the Western Ghats. In contrast, middle level cloud fraction shows large values (~ 40 %) compared to high cloud fraction over Coastal regions of Myanmar, Thailand and Cambodia as well as over the Western Ghats over Indian peninsula. From Fig. 4, it can be observed that the three negative NETCRF regions (in Fig. 3) are characterized by large values of high and middle level cloud fraction compared to rest of the Indian region. High cloud fraction dominates the cloudiness over the Bay of Bengal region, while upper middle and high cloud fractions dominate the land NETCRF regimes. However, differentiating the negative NETCRF regimes merely based on the cloud cover data alone is inadequate. In order to have a quantitative understanding regarding parameters influencing the three negative NETCRF regimes, seasonal mean variation of various cloud and environmental parameters over these regions are examined.

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The seasonal mean values of cloud parameters (cloud fraction, particle size, optical depth) and environmental variables (rain rate, Free tropospheric humidity (FTH), precipitable water) over the three negative NETCRF regions are calculated for the summer monsoon season of 2002–2005 and are shown in Table 4. From Table 4, it can be seen that high and upper middle cloud fraction dominates the total cloudiness over the three regions. High cloud amount ($\sim 51\%$) and optical depth (~ 14) are largest over the Bay of Bengal while upper middle cloud amount shows large values ($\sim 34\%$ and 43%) compared to high cloud amount over the land areas. Amount of lower level (lower middle and low) clouds are below 10% over the Bay of Bengal while, it shows values between 10 – 15% over land regions. Earlier studies showed that high clouds account for $\sim 65\%$ of all cloud grids observed over the Indian monsoon region during the monsoon season (Tang and Chen, 2006; Meenu et al., 2007). Relatively lower values of low level clouds (compared to high and middle level clouds) are observed over the three regions, which could be due to underestimation of low level clouds by MODIS and geostationary instruments over the Indian monsoon region (Tang and Chen, 2006). The mean ice and water cloud particle size for the three regions show similar values while precipitable water vapor values (from GMAO GEOS database) show a variation between 48 to 58 mm for the three regions. From this analysis, it is not easy to comprehend the influence of various parameters on the negative NETCRF over these regions. In order to obtain a better understanding regarding the influence of these parameters on the negative NETCRF over the three regions, TOA flux and CRF values for the three regions are computed using RRTM simulations. The regional mean values of various cloud and environmental variables estimated for the three regions are used as input in the model. Using the cloud cover and cloud particle size data from CERES, altitude profile of cloud cover and SSA for each region is developed employing the parameterization schemes described in Sect. 3. TOA flux values estimated using the RRTM simulations for the three regions are used to compute the NETCRF.

5 Validation of TOA flux and CRF: model vs. observations

The seasonal mean TOA flux estimated for the three negative NETCRF regions from CERES, ISCCP-FD and RRTM simulations are shown in Table 5. The ISCCP FD TOA flux data belonging to the study period is used in the estimation of seasonal mean TOA flux and CRF. In general, TOA flux from CERES observations and RRTM simulations are very much in agreement with each other. However, TOA flux derived from ISCCP FD flux data shows consistently larger values ($\sim 5\text{--}10\text{ W m}^{-2}$) compared to CERES and RRTM estimations. For example, CERES TOA clear sky SW flux values (in W m^{-2}) are 46.8, 63.4, and 61.9 over the Bay of Bengal, Myanmar and the Western Ghats respectively while the same from ISCCP FD (in W m^{-2}) are 55.4, 71 and 68.6, indicating an overestimation (of the order of $7\text{--}9\text{ W m}^{-2}$) by the ISCCP FD data. Significant variations between the CERES and RRTM derived fluxes are observed only in the case of total sky flux estimated over Myanmar and Western Ghats region. Using the TOA flux values (presented in Table 5), SW, LW and NET CRF values for the three regions are calculated and presented in Table 6. From Table 6, it can be seen that the CERES and RRTM derived CRF values are very much in agreement over the Bay of Bengal region whereas the ISCCP FD derived values show considerable difference ($4\text{--}8\text{ W m}^{-2}$). Over the land regions, both the RRTM and ISCCP derived SW and LW CRF values show substantial variation from CERES derived values. Still, RRTM derived SW and LW CRF values are much closer to the CERES derived values compared that from ISCCP FD. The NETCRF estimated from the CERES and RRTM shows very good agreement (within the uncertainty limit of CERES observations) while the ISCCP values are well outside the uncertainty limit of CERES observation.

The NETCRF estimated from ISCCP FD data over the three regions shows consistently large values (in magnitude) compared to CERES and RRTM derived values. This indicates that the cloud vertical model used in the estimation of TOA flux (RRTM) for the three negative NETCRF regimes are more accurate in representing the cloud overlap and altitude structure compared to the cloud vertical structure (CVS) scheme

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



used in the ISCCP FD flux estimation. However, it can be argued that the ISCCP CVS scheme is a general overlap scheme applicable for a broad range of meteorological conditions, while the present cloud overlap scheme is tailored for a specific convective cloud condition. The disparity in SW and LW CRF values computed by the RRTM and CERES over the land regimes (Myanmar and the Western Ghats) can be attributed to the underlying uncertainty associated with the parameterization of cloud vertical structure and its microphysical properties. After successfully simulating the negative NETCRF regimes over the Indian region using RRTM, the next step is to identify the parameters that control the negative NETCRF over these regions. This is carried out by analyzing the sensitivity of cloud radiative forcing to various cloud and environmental variables (using RRTM) and explained in the subsequent sections.

5.1 Sensitivity calculations

In this section, sensitivity of CRF to various cloud macro/micro physical properties and environmental variables are analyzed by studying their relative contribution to the NETCRF. In the earlier section, it was shown that negative NETCRF over the three regions could be modeled with good accuracy using RRTM and cloud parameterization schemes. Using the same simulation methods, it is possible to quantify the dependence of NETCRF on various cloud micro/macro physical and environmental variables over the three regions. This also provides an opportunity to test the veracity of theoretical hypothesis propagated by various investigators on the occurrence of negative NETCRF over tropical convective regions. In this analysis, influence of cloud amount, cloud particle size, single scattering albedo and atmospheric water vapor on the negative NETCRF over the three regions are examined.

5.1.1 Influence of cloud macro-physical properties on CRF

In the first simulation, influence of cloud top altitude on the LW and SW CRF values are examined for the three regions. In this analysis, TOA flux values are computed using

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



RRTM for the three regions by varying the cloud top altitude in the respective cloud vertical model from 125 hPa to 400 hPa in several steps while keeping all other input parameters constant. The resultant LW and SW CRF for different cloud top pressures over the Bay of Bengal (thick line) and Myanmar (dashed line) region are estimated and shown in Fig. 5. From Fig. 5, it can be seen that LWCRF decreases monotonically with increase in the cloud top pressure (from ~ 100 and 80 W m^{-2} respectively to ~ 45 and 40 W m^{-2} , respectively) for the two regions while the SWCRF shows very little variation ($< 5 \text{ W m}^{-2}$). This indicates that change in cloud top altitude of high clouds cause relatively large variation in LWCRF ($\sim 50 \text{ W m}^{-2}$) compared to SWCRF, which results in causing an imbalance between the two. The sharp wedge observed in the SWCRF variation in Fig. 5 is due to the change in cloud cover type from high level to middle level. These results are in agreement with that of Kiehl (1994) who showed that decrease (increase) in cloud top altitude causes system to shift towards a negative (positive) NETCRF regime. However, seasonal mean cloud top altitude (high clouds) over the Indian region during monsoon season shows a variation between 180–240 hPa. For such variation in cloud top altitude, corresponding variation in LWCRF observed for the three regions are less than 13 W m^{-2} . This shows that small variation in cloud top altitude of high clouds cannot significantly influence the magnitude of NETCRF over the Indian region.

In the second sensitivity analysis, we try to quantify the competing influence of cloud top and cloud amount on the LWCRF and NETCRF. In this analysis, high cloud amount in the cloud vertical model representing the Bay of Bengal regime is varied from its original value of 50 % to 20 % in steps of 10 %, keeping cloud amount of other groups in the cloud vertical model same. By stepwise reduction in high cloud amount from cloud vertical model, fraction of upper middle level clouds exposed to the TOA (and to the incoming solar incoming radiation) increases from 12 % to 42 %. Running RRTM with this modified cloud vertical profile shows a decrease in LWCRF from 78 W m^{-2} to 66 W m^{-2} while the SWCRF show very little variation since the total cloud cover remains the same (top level clouds are sliced off while bottom level clouds remain intact).

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Impact of this reduction in high cloud amount is to increase the absolute magnitude of the NETCRF (by 12 W m^{-2}) and shift NETCRF the region towards a stronger negative regime due the decrease of LWCRF. Similar analysis was performed for the Myanmar and the Western Ghats region where it showed similar result as that of the Bay of Bengal region. Cess et al. (2001) observed that change in cloud vertical structure associated with El Nino over the tropical Pacific Ocean tend to cause substantial radiative cooling. It was observed that CRF over Pacific warm pool is partially governed by high and middle-level clouds during the El Nino year compared to high level clouds during normal years. This analysis indicates that in a multi layered cloud system, both cloud vertical structure and cloud amount influences the NETCRF, even though magnitude of variation in NETCRF is not significantly large in this case.

In the third test, influence of low levels clouds on the NETCRF is examined over the three regions while keeping all other parameters in the cloud vertical model constant. In this analysis, un-observed portions (visible from TOA) of the low levels clouds are removed from the cloud vertical profile and LWCRF and SWCRF values are recomputed using the modified cloud vertical model. The modified LW and SW CRF values estimated from the analysis are presented in Table 7. From Table 7, it can be seen that the low level cloud cover over these regions has very little influence in controlling the observed LWCRF and SWCRF compared to high level clouds. The SWCRF over the Western Ghats shows a decrease ($\sim 7 \text{ W m}^{-2}$) when the modified cloud vertical profile (low level clouds removed) is used. For the Bay of Bengal and Myanmar region, the maximum decrease in SWCRF is of the order of $< 3 \text{ W m}^{-2}$. Corresponding variation in the LWCRF is considerably small compared to that of SWCRF. Large low level cloud fraction ($\sim 14\%$) was observed over the Western Ghats while it is $< 6\%$ for the other two regions. Present analysis shows that low level cloud amount has little influence in modulating the NETCRF values over the three study regions.

5.1.2 Influence of cloud microphysical properties on CRF

In the present analysis, influence of cloud single scattering albedo and ice cloud particle size on NETCRF is investigated using RRTM. The layer averaged cloud particle size (from SRBAVG2 GEO data) for ice clouds over the Indian region show variation of effective particle size between the range 18–30 μm . Cloud ice particle size is directly related to the cloud single scattering properties (Eq. 11), which in turn modulates the SWCRF and NETCRF of the cloud system. In the present analysis, an attempt is made to quantify the sensitivity of SWCRF to the ice cloud particle size. This is done by computing the SWCRF over the three regions by varying the ice particle size of high and upper middle level clouds (there by varying the SSA) in the cloud vertical model while keeping all other parameters constant. The ice particle size is varied from 20 to 40 μm and SWCRF values are computed each time using RRTM by incorporating the modified SSA in the input cloud model. SWCRF estimated from the model simulations are presented in Table 8. From Table 8, it can be seen that the SWCRF values show a monotonic decrease with increasing ice particle size for the three regions. A maximum decrease in SWCRF value of $\sim 10 \text{ W m}^{-2}$ for an increase in ice particle size from 20 to 40 μm is observed over the Bay of Bengal region while it is much less over the other two regions. The study indicates that increase in particle size leads to moderate decrease in SWCRF and NETCRF over the study region. Studies have shown changes in cloud particle (ice) size can modify the net radiative forcing of cirrus clouds to a cooling or warming regime depending on the direction of change (Zhang et al., 1999). However, it is seems unlikely that small variation in cloud ice particle size (between 18–30 μm) over the Indian region alone can bring about the negative NETCRF values observed over the region.

5.1.3 Influence of atmospheric water vapor on CRF

In this section, sensitivity analysis carried out using RRTM to understand the influence of atmospheric water vapor on the LWCRF over the three negative NETCRF regions

Negative cloud forcing

B. V. Thampi and R. Roca

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



are explained. Studies have shown that atmospheric water vapor can significantly influence the LW forcing by reducing the clear sky TOA flux (Sohn and Schmetz, 2004; Sohn and Bennartz, 2008). Water vapor being an important greenhouse gas absorbs the atmospheric LW radiation and decreases the LWCRF (while having little effect on SWCRF). Roca et al. (2004) proposed that damping of LWCRF by large amount of water vapor present in the atmosphere over the Bay of Bengal region during the summer monsoon season could be a reason for the observed negative NETCRF. In this analysis, CRF over the three negative NETCRF regions are simulated by varying the PW vapor content in the atmosphere from their original values (between 48–58 mm) while keeping all other parameters in model simulation constant. This is achieved by varying the relative humidity (thereby water vapor mixing ratio) of free troposphere in the model atmosphere while keeping the boundary layer and stratospheric RH constant. This results in the formation of several model atmospheric profiles with distinctly different PW values (between 33 to 68 mm). Using this methodology, variation of TOA LW flux and LWCRF with PW for the three regions is examined.

Variation of TOA LW flux (clear and total sky) and LWCRF with PW for the Bay of Bengal region is shown in Fig. 6. In general, clear and total sky LW flux shows a monotonic decrease with increase in PW. TOA LW clear sky flux over the Bay of Bengal region shows a decrease of $\sim 45 \text{ W m}^{-2}$ for an increase in PW value from 33 to 68 mm while the total sky LW flux show a decrease of $\sim 20 \text{ W m}^{-2}$. The LWCRF also show considerable decrease over the Bay of Bengal region from 106 W m^{-2} to 74 W m^{-2} (decrease of $\sim 32 \text{ W m}^{-2}$) for a corresponding increase in PW from 33 mm to 68 mm. Over Myanmar and the Western Ghats, LWCRF shows a decrease of ~ 30 and $\sim 25 \text{ W m}^{-2}$ respectively, for a similar variation in PW from 33 mm to 68 mm. Sohn et al. (2006) demonstrated that water vapor in the upper troposphere can contribute 12 W m^{-2} to the LWCRF over convectively active tropical regions. This analysis shows that atmospheric water vapor can cause a relatively larger variation in LWCRF (and in NETCRF) compared to other variables discussed in earlier sections. For a through understanding, seasonal mean variation in PW over the Indian region during different

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



seasons is also examined. Seasonal mean variation of precipitable water vapor (from the CERES SRBAVG GEO/GMAO GEOS product) over the Indian region during the summer and winter months are estimated and shown in Fig. 7. From Fig. 7a and b, it can be seen that average value of PW observed over the Indian region is between 40 and 60 mm (summer months), while it is between ~ 20 –30 mm during the winter months (DJF). This indicates that between dry (winter) and wet (summer) seasons, PW content in the atmosphere over the Indian region increases by ~ 25 –35 mm. RRTM simulations shows that such increase in PW can cause a substantial decrease in clear sky TOA flux (~ 30 –40 W m^{-2}) and LWCRF (~ 20 –30 W m^{-2}). From Fig. 6, it can be seen that LWCRF over Bay of Bengal increase by $\sim 23 \text{ W m}^{-2}$ for a decrease in PW content from 58 mm to 30 mm. Over the two land regimes, a similar increase in LWCRF is observed though magnitude of increase in LWCRF is lower than that observed over the Bay of Bengal region. From this analysis, it can be seen that the atmospheric water content over these regions significantly modifies the observed LWCRF and NETCRF values. Hence, relatively large amount of water vapor in the atmosphere over the three negative NETCRF regions during the summer monsoon season is a major factor controlling the imbalance between SWCRF and LWCRF (and the negative NETCRF regimes).

6 Inter-comparison of the oceanic NETCRF regimes: Bay of Bengal vs. West Pacific

We now consider two specific oceanic convective regions for a better understanding of the negative and near zero NETCRF regimes; the Bay of Bengal (10 – 22°N ; 85 – 100°E) during the Asian monsoon season (represent negative regime) and the West Pacific (10°S – 10°N ; 140°E – 180°E) during the March–April period (near zero regime). The two regions share similar surface properties (ocean surface type) with almost similar sea surface temperature (SST). Figure 8 presents the mean variation of CERES TOA NETCRF over the West Pacific during the March–April period for the years 2002–2005. From Fig. 8, it can be seen that the regional variation of NETCRF over the West

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Pacific during this period lies mainly in the range of $\pm 20 \text{ W m}^{-2}$. Negative NETCRF values ($\sim 20 \text{ W m}^{-2}$) are observed over small areas close to the Indonesian islands while rest of the region show positive or near zero NETCRF values. A similar regional variation of NETCRF over the Bay of Bengal (Fig. 3) region during the monsoon season shows negative NETCRF values ranging from -10 to -70 W m^{-2} . For a more quantitative assessment, mean TOA flux and CRF over the two regions are estimated and presented in Table 9. In general, the TOA SW (clear and total) flux values are largest over the Bay of Bengal region compared to the West Pacific while the reverse is true for TOA LW fluxes. Another interesting point here is the difference in clear sky TOA LW flux between two regions; relatively lower TOA LW clear sky flux (276 W m^{-2}) values are observed over the Bay of Bengal compared to that over the West Pacific (285 W m^{-2}). While LWCRF and SWCRF values over Bay of Bengal region shows large imbalance (resulting in a NETCRF value $\sim -30 \text{ W m}^{-2}$), that over the West Pacific shows a near balance between LWCRF and SWCRF (65 and -67 W m^{-2} , respectively) leading to near zero NETCRF over the West Pacific. The disparity in the estimated CRF values observed over the two oceanic regions are evident in Table 9.

Table 10 presents the regional mean cloud fraction and cloud optical depth estimated using CERES data over the Bay of Bengal and the West Pacific. From Table 10, it can be seen that high level cloud fraction ($\sim 51\%$) observed over the Bay of Bengal is relatively large compared to that observed over the West Pacific ($\sim 38\%$) while, lower level cloud types show almost similar variation over the two regions. A comparison of cloud optical depth and cloud top altitude between the various cloud groups also show similar variations. This indicates that cloud properties over the two convective regions show a lot of similarities except for the high level cloud fraction. However, the interesting question here is “whether the variation observed in the cloud amount (13% and 5% in high and upper middle cloud cover respectively between these two regions) alone can cause the NETCRF to shift between near zero and negative values”? Rajeevan and Sreenivasan (2000) postulated that large high cloud amount observed over the Bay of Bengal region is the reason behind the large negative NETCRF compared to rest of

tropical convective regions. However, RRTM simulations incorporating the mean cloud and environmental variables over the West Pacific region show that such variability in cloud cover alone is not enough to drive the system from a near zero to large negative NETCRF regime.

The other important parameters that can influence the NETCRF over the two regions are cloud microphysical properties (cloud particle size, shape etc) and atmospheric water content. Comparison of cloud particle size ($\sim 25 \mu\text{m}$ for ice and $\sim 12 \mu\text{m}$ for water clouds, respectively) over the two regions shows very little variation. The other variable that can influence the NETCRF over the two regions is the atmospheric water vapor content. In Sect. 5.1.3, influence of atmospheric water vapor on the TOA flux and CRF over the Indian region was discussed. A similar analysis is carried out to quantify the influence of atmospheric water vapor on the NETCRF over the West Pacific. For this purpose, monthly mean atmospheric water vapor content over the two regions from special sensor microwave/imager (SSM/I) data (Wentz, 1997) was estimated for the April (West Pacific) and July (Bay of Bengal) months (2002–2005) and is shown in Fig. 9. The atmospheric water vapor content over the Bay of Bengal during the July month is extremely large ($> 58 \text{ mm}$) compared to that observed over the West Pacific region ($\sim 45\text{--}55 \text{ mm}$) during April. Over the negative NETCRF regions ($< -30 \text{ W m}^{-2}$) of Bay of Bengal, the PW reaches as high as $\sim 68 \text{ mm}$. The water vapor information over the land is not available from SSM/I data. However, from Fig. 9, it can be seen that the water vapor amount progressively increases towards land area (Myanmar coast) over the Bay of Bengal region. Large values of atmospheric water vapor ($> 65 \text{ mm}$) are observed over the Bay of Bengal region close to Myanmar coast where the NETCRF values also show negative values ($< -30 \text{ W m}^{-2}$). Compared to the Bay of Bengal region, the water vapor loading over the West Pacific usually lies within $48\text{--}58 \text{ mm}$ during the April month. Inter-annual variation of atmospheric water vapor content over the Bay of Bengal region shows that precipitable water vapor amount reach as high as $\sim 70 \text{ mm}$ over the core negative NETCRF regions during the summer monsoon seasons, while it hardly goes beyond 60 mm over the West Pacific during March–April. The redistribu-

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tion of water vapor associated with convection results in increased moistening of the upper troposphere over the Indian region, which contribute significantly to the damping of LWCRF and creating an imbalance with SWCRF. Magnitude of such damping in LWCRF over the West Pacific is relatively small compared to the Bay of Bengal region.

- 5 The increase in atmospheric water vapor content and high cloud amount could be the reason behind negative NETCRF values being observed over the Bay of Bengal region compared to near zero NETCRF values observed over the West Pacific region.

7 Summary and discussion

The net cloud radiative forcing over the Indian land mass and adjacent oceanic regions during the Asian summer monsoon season of 2002–2005 were investigated using CERES observations and RRTM simulations. The seasonal mean variation of NETCRF over the Indian region during the Asian summer monsoon season showed the existence of three strong negative NETCRF regimes; Bay of Bengal ($\sim -31 \text{ W m}^{-2}$), Myanmar ($\sim -37 \text{ W m}^{-2}$) and the Western Ghats ($\sim -31 \text{ W m}^{-2}$), respectively. Using RRTM and cloud vertical model, the TOA flux and CRF over the negative NETCRF regions during the summer monsoon season was studied. Using the cloud vertical model as input, NETCRF for the three regions were estimated using RRTM. The NETCRF values calculated from RRTM simulations found to agree well with CERES observations, while that from ISCCP FD data showed large differences. Sensitivity of the negative NETCRF values to various cloud micro/macro physical and environmental variables were carried out using model simulations. Sensitivity of ice particle size to the NETCRF was evaluated by varying the ice particle radius (from 20 to 40 μm) in the parameterization of SSA, which produced a maximum variation of $\sim 10 \text{ W m}^{-2}$ on the NETCRF values. Decrease in cloud particle size found to increase the SWCRF with very little variation in LWCRF. However, variations in CRF due to changes in cloud particle size alone is not sufficient enough to cause the formation of negative NETCRF regimes observed over the Indian region.

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Studies showed that low level cloud fraction has very little influence in controlling the NETCRF values over the three negative NETCRF regions. RRTM simulations of LWCRF employing model atmosphere with varying PW values showed that atmospheric water vapor content significantly influence the NETCRF values observed over the region. A decrease in LWCRF by $\sim 25\text{--}35\text{ W m}^{-2}$ is observed for an increase of PW from $25\text{--}35\text{ mm}$ over the negative NETCRF regions. The precipitable water amount is found to be higher ($> 55\text{ mm}$) over the Bay of Bengal and Myanmar region compared to other locations over the Indian region during the monsoon season. Similarly, the clear sky TOA LW flux over Bay of Bengal and Myanmar region was found to be significantly smaller ($\sim 276\text{ W m}^{-2}$) during the summer monsoon season compared to winter months ($\sim 300\text{ W m}^{-2}$). This increase in atmospheric water vapor associated with the convective activity over the Indian region during the monsoon season reduces the TOA LW clear sky flux and LWCRF. Influence of water vapor loading is largest over the North Bay of Bengal region close to the Myanmar coast where the total cloud amount is also largest. This indicates that the combined effect of large amount of high level clouds and increased atmospheric water vapor loading over the Indian region significantly influence the imbalance between LWCRF and SWCRF. Inter comparison between CRF values observed over the Bay of Bengal and the West Pacific also indicates the importance of atmospheric water vapor in controlling the magnitude of NETCRF values. Precipitable water vapor content over the North Bay of Bengal region reaches as high as $\sim 70\text{ mm}$ during the monsoon months while it hardly reaches 60 mm over the West Pacific. The damping of LWCRF caused by the increased atmospheric water vapor loading and presence of large amount of high clouds over the Indian region during the summer monsoon season creates an imbalance between SWCRF and LWCRF and is accountable for the observed negative NETCRF values over the region.

Acknowledgements. The TOA flux data used in the study were obtained from the NASA Langley Research Center EOSDIS Distributed Active Archive Center. The ISCCP FD data were obtained from the International Satellite Cloud Climatology (<http://isccp.giss.nasa.gov>). Authors thank the SSM/I for providing the atmospheric water vapor data used in the study.

Technical and data support from the Climserv/LMD team is fully appreciated. Authors thank Eli Mlawer and the Atmospheric and Environmental Research (AER) team for providing the RRTMG model (<http://rtweb.aer.com/>). B. V. Thampi acknowledges CNES, France for providing the post-doc research fellowship. Authors acknowledge the support and help of colleagues at LMD/UPMC, France.



The publication of this article is financed by CNRS-INSU.

References

- 10 Balachandran, S. and Rajeevan, M.: Sensitivity of surface radiation budget to clouds over the Asian monsoon region, *J. Earth Syst. Sci.*, 116, 159–169, 2007.
- Barker, H. W., Stephens, G. L., Partain, P. T., Bergman, J. B., Bonnel, B., Campana, K., Clothiaux, E. E., Clough, S., Cusack, S., Delamere, J., Edwards, J., Evans, K. F., Fouquart, Y., Freidenreich, S., Galin, V., Hou, Y., Kato, S., Li, J., Mlawer, E., Morcrette, J.-J., O'Hirok, W., Räsänen, P., Ramaswamy, V., Ritter, B., Rozanov, E., Schlesinger, M., Shibata, K., Sporyshev, P., Sun, Z., Wendisch, M., Wood, N., and Yang, F.: Assessing 1-D atmospheric solar radiative transfer models: interpretation and handling of unresolved clouds, *J. Climate*, 16, 2676–2699, 2003.
- 15 Bergman, J. W. and Hendon, H. H.: Calculating monthly radiative fluxes and heating rates from monthly cloud observations, *J. Atmos. Sci.*, 55, 3471–3492, 1998.
- Bodas-Salcedo, A., Webb, M. J., Bony, S., Chepfer, H., Dufresne, J.-L., Klein, S. A., Zhang, Y., Marchand, R., Haynes, J. M., Pincus, R., and John, V. O.: COSP: satellite simulation software for model assessment, *B. Am. Meteorol. Soc.*, 92, 1023–1043, 2011.
- 25 Cess, R. D., Zhang, M., Wielicki, B. A., Young, D. F., Zhou, X. L., and Nikitenko, Y.: The influence of the 1998 El Niño upon cloud-radiative forcing over the pacific warm pool, *J. Climate*, 14, 2129–2137, 2001.

- Chambon, P., Jobard, I., Roca, R., and Viltard, N.: An investigation of the error budget of tropical rainfall accumulation derived from merged passive microwave and infrared satellite measurements, *Q. J. Roy. Meteor. Soc.*, 139, 879–893, doi:10.1002/qj.1907, 2011.
- Charlock, T. P. and Ramanathan, V.: The albedo field and cloud radiative forcing produced by a general circulation model with internally generated cloud optics, *J. Atmos. Sci.*, 42, 1408–1429, 1985.
- Chou, M.-D., Suarez, M. J., Ho, C.-H., Yan, M. M.-H., and Lee, K.-T.: Parameterizations for cloud overlapping and shortwave single-scattering properties for use in general circulation and cloud ensemble models, *J. Climate*, 11, 202–214, 1998.
- Clough, S. A., Shephard, M. W., Mlawer, E. J., Delamere, J. S., Iacono, M. J., Cady-Pereira, K., Boukabara, S., and Brown, P. D.: Atmospheric radiative transfer modeling: a summary of the AER codes, *J. Quant. Spectrosc. Ra.*, 91, 233–244, 2005.
- Cole, J., Barker, W. H., Loeb, N. G., and Salzen, K.: Assessing simulated clouds and radiative fluxes using properties of clouds whose tops are exposed to space, *J. Climate*, 24, 2715–2727, 2011.
- Collins, W. D.: Parameterization of generalized cloud overlap for radiative calculations in general circulation models, *J. Atmos. Sci.*, 58, 3224–3242, 2001.
- Dessler, A. E. and Yang, P.: The distribution of tropical thin cirrus clouds inferred from Terra MODIS data, *J. Climate*, 16, 1241–1247, 2003.
- Devasthale, A. and Fueglistaler, S.: A climatological perspective of deep convection penetrating the TTL during the Indian summer monsoon from the AVHRR and MODIS instruments, *Atmos. Chem. Phys.*, 10, 4573–4582, doi:10.5194/acp-10-4573-2010, 2010.
- Ellingson, R. G. and Fouquart, Y.: The intercomparison of radiation codes in climate models: an overview, *J. Geophys. Res.*, 96, 8925–8927, 1991.
- Ellingson, R. G. and Wiscombe, W. J.: The Spectral Radiance Experiment (SPECTRE): project description and sample results, *B. Am. Meteorol. Soc.*, 77, 1967–1985, 1996.
- Evans, J. S., Cooper, D. W., and Kinney, P.: On the propagation of error in air pollution measurements, *Environ. Monit. Assess.*, 4, 139–153, 1984.
- Fouquart, Y.: Radiation in boundary layer clouds, in: Report of the JSC/CAS Workshop on Modelling of Cloud-Topped Boundary Layer, Fort Collins, Colorado, USA, 22–26 April, Appendix D, 40 pp., 1985.
- Fouquart, Y. and Bonnel, B.: Computations of solar heating of the earth's atmosphere: a new parameterization, *Beitr. Phys. Atmos.*, 53, 35–62, 1980.

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Fu, Q.: An accurate parameterization of the solar radiative properties of cirrus clouds for climate models, *J. Climate*, 9, 2058–2082, 1996.
- Futyan, J. M., Russell, J. E., and Harries, J. E.: Cloud radiative forcing in Pacific, African, and Atlantic tropical convective regions, *J. Climate*, 17, 3192–3202, 2004.
- 5 Geleyn, J.-F. and Hollingsworth, A.: An economical analytical method for the computation of the interaction between scattering and line absorption of radiation, *Beitr. Phys. Atmos.*, 52, 1–16, 1979.
- Hahn, C. J., Rossow, W. B., and Warren, S. G.: ISCCP cloud properties associated with standard cloud types identified in individual surface observations, *J. Climate*, 14, 11–28, 2001.
- 10 Harrison, E. F., Minnis, P., Barkstrom, B. R., Ramanathan, V., Cess, R. D., and Gibson, G. G.: Seasonal variation of cloud radiative forcing derived from the Earth Radiation Budget Experiment, *J. Geophys. Res.*, 95, 18687–18703, 1990.
- Hartmann, D. L., Moy, L. A., and Fu, Q.: Tropical convection and the energy balance at the top of the atmosphere, *J. Climate*, 14, 4495–4511, 2001.
- 15 Hu, Y. X. and Stamnes, K.: An accurate parameterization of the radiative properties of water clouds suitable for use in climate models, *J. Climate*, 6, 728–742, 1993.
- Iacono, M. J., Mlawer, E. J., Clough, S. A., and Morcrette, J.-J.: Impact of an improved longwave radiation model, RRTM, on the energy budget and thermodynamic properties of the NCAR community climate model, CCM3, *J. Geophys. Res.*, 105, 14873–14890, doi:10.1029/2000JD900091, 2000.
- 20 Jensen, E. J., Kinne, S., and Toon, O. B.: Tropical cirrus cloud radiative forcing: sensitivity studies, *Geophys. Res. Lett.*, 21, 2023–2026, 1994.
- Key, J. R., and Schweiger, A. J.: Tools for atmospheric radiative transfer: Streamer and FluxNet, *Comput. Geosci.*, 24, 443–451, doi:10.1016/S0098-3004(97)00130-1, 1998.
- 25 Key, J. R., Yang, P., Baum, B. A., and Nasiri, S. L.: Parameterization of shortwave ice cloud optical properties for various particle habits, *J. Geophys. Res.*, 107, AAC 7-1–AAC 7-10, doi:10.1029/2001JD000742, 2002.
- Kiehl, J. T.: On the observed near cancellation between longwave and shortwave cloud forcing in tropical regions, *J. Climate*, 7, 559–565, 1994.
- 30 Kiehl, J. T. and Ramanathan, V.: Comparison of cloud forcing derived from Earth Radiation Budget Experiment with that simulated by the NCAR community climate model, *J. Geophys. Res.*, 95, 1679–1698, 1990.

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Klein, S. A. and Jakob, C.: Validation and sensitivities of frontal clouds simulated by the ECMWF model, *Mon. Weather Rev.*, 127, 2514–2531, 1999.
- Liou, K.-N.: Influence of cirrus clouds on weather and climate processes: a global perspective, *Mon. Weather Rev.*, 114, 1167–1199, 1986.
- 5 Liou, K. N. and Ou, S. C.: The role of cloud microphysical processes in climate: an assessment from a one-dimensional perspective, *J. Geophys. Res.*, 94, 8599–8607, 1989.
- Lo, E.: Gaussian error propagation applied to ecological data: post-ice-storm-downed woody biomass, *Ecol. Monogr.*, 75, 451–466, 2005.
- Loeb, N. G., Kato, S., Loukachine, K., and Manalo-Smith, N.: Angular distribution models for
 10 top-of-atmosphere radiative flux estimation from the clouds and the Earth's radiant energy system instrument on the Terra satellite, Part I: Methodology, *J. Atmos. Ocean. Tech.*, 22, 338–351, 2005.
- Loeb, N. G., Wielicki, B. A., Doelling, D. R., Smith, G. L., Keyes, D. F., Kato, S., Manalo-Smith, N., and Wong, T.: Toward optimal closure of the earth's top-of-atmosphere radiation
 15 budget, *J. Climate*, 22, 748–766, 2009a.
- Loeb, N. G., Wielicki, B. A., Wong, T., and Parker, P. A.: Impact of data gaps on satellite broadband radiation records, *J. Geophys. Res.*, 114, D11109, doi:10.1029/2008JD011183, 2009b.
- Meenu, S., Rajeev, K., Parameswaran, K., and Suresh Raju, C.: Characteristics of the double intertropical convergence zone over the tropical Indian Ocean, *J. Geophys. Res.*, 112,
 20 D11106, doi:10.1029/2006JD007950, 2007.
- Meenu, S., Rajeev, K., Parameswaran, K., and Nair, A. K. M.: Regional distribution of deep clouds and cloud top altitudes over the Indian subcontinent and the surrounding oceans, *J. Geophys. Res.*, 115, D05205, doi:10.1029/2009JD011802, 2010.
- Menzel, W. P., Frey, R. A., Zhang, H., Wylie, D. P., Moeller, C. C., Holz, R. E., Maddux, B.,
 25 Baum, B. A., Strabala, K. I., and Gumley, L. E.: MODIS global cloud top pressure and amount estimation: algorithm description and results, *J. Appl. Meteorol. Clim.*, 47, 1175–1198, doi:10.1175/2007JAMC1705.1, 2008.
- Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J., and Clough, S. A.: RRTM, a validated correlated- k model for the longwave, *J. Geophys. Res.*, 102, 16663–16682, 1997.
- 30 Morcrette, J.-J., Barker, H. W., Cole, J., Iacono, M., and Pincus, R.: Impact of a new radiation package, mcrad, in the ecmwf integrated forecasting system, *Mon. Weather Rev.*, 136, 4773–4798, 2008.

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Pai, D. S. and Rajeevan, M.: Clouds and cloud radiative forcing over tropical Indian ocean and their relationship with sea surface temperature, *Curr. Sci. India*, 75, 372–381, 1998.
- Patil, S. D. and Yadav, R. K.: Large-scale changes in the cloud radiative forcing over the Indian region, *Atmos. Environ.*, 39, 4609–4618, 2005.
- 5 Pavlaklis, K. G., Hatzianastassiou, N., Matsoukas, C., Fotiadi, A., and Vardavas, I.: ENSO surface shortwave radiation forcing over the tropical Pacific, *Atmos. Chem. Phys.*, 8, 5565–5577, doi:10.5194/acp-8-5565-2008, 2008.
- Pincus, R., Barker, H. W., and Morcrette, J.-J.: A fast, flexible, approximate technique for computing radiative transfer in inhomogeneous cloud fields, *J. Geophys. Res.*, 108, 4376, doi:10.1029/2002JD003322, 2003.
- 10 Platt, C. M. R.: A parameterization of the visible extinction coefficient of ice clouds in terms of the ice/water content, *J. Atmos. Sci.*, 54, 2083–2098, 1997.
- Räsänen, P.: Parameterization of water and ice cloud near-infrared single-scattering co-albedo in broadband radiation schemes, *J. Atmos. Sci.*, 56, 626–641, 1999.
- 15 Rajeevan, M. and Srinivasan, J.: Net cloud radiative forcing at the top of the atmosphere in the Asian monsoon region, *J. Climate*, 13, 650–657, 2000.
- Ramanathan, V., Cess, R. D., Harrison, E. F., Minnis, P., Barkstrom, B. R., Ahmad, E., and Hartmann, D.: Cloud-radiative forcing and climate: results from the Earth Radiation Budget Experiment, *Science*, 243, 57–63, 1989.
- 20 Remer, L. A., Kaufman, Y. J., Tanré, D., Mattoo, S., Chu, D. A., Martins, J. V., Li, R.-R., Ichoku, C., Levy, R. C., Kleidman, R. G., Eck, T. F., Vermote, E., and Holben, B. N.: The MODIS aerosol algorithm, products, and validation, *J. Atmos. Sci.*, 62, 947–973, 2005.
- Roca, R., Louvet, S., Picon, L., and Desbois, M.: A study of convective systems, water vapor and top of the atmosphere cloud radiative forcing over the Indian Ocean using INSAT-1B and ERBE data, *Meteorol. Atmos. Phys.*, 90, 49–65, 2004.
- 25 Rossow, W. B. and Zhang, Y. C.: Calculation of surface and top of atmosphere radiative fluxes from physical quantities based on ISCCP datasets. 2. Validation and first results, *J. Geophys. Res.*, 100, 1167–1197, 1995.
- Rossow, W. B., Zhang, Y., and Wang, J.: A statistical model of cloud vertical structure based on reconciling cloud layer amounts inferred from satellites and radiosonde humidity profiles, *J. Climate*, 18, 3587–3605, 2005.
- 30

- Schmidt, E. O., Arduini, R. F., Wielicki, B. A., Stone, R. S., and Tsay, S.-C.: Considerations for modeling thin cirrus effects via brightness temperature differences, *J. Appl. Meteorol.*, 34, 447–459, 1995.
- Slingo, A.: A GCM parameterization for the shortwave radiative properties of water clouds, *J. Atmos. Sci.*, 46, 1419–1427, 1989.
- Slingo, A. and Schrecker, H. M.: On the shortwave radiative properties of stratiform water clouds, *Q. J. Roy. Meteor. Soc.*, 108, 407–426, 1982.
- Smith, G. L., Mlynczak, P. E., and Potter, G. L.: A technique using principal component analysis to compare seasonal cycles of earth radiation from CERES and model computations, *J. Geophys. Res.*, 117, D09116, doi:10.1029/2011JD017343, 2012.
- Sohn, B.-J. and Bennartz, R.: Contribution of water vapor to observational estimates of long-wave cloud radiative forcing, *J. Geophys. Res.*, 113, D20107, doi:10.1029/2008JD010053, 2008.
- Sohn, B.-J. and Schmetz, J.: Water vapor-induced OLR variations associated with high cloud changes over the tropics: a study from Meteosat-5 observations, *J. Climate*, 17, 1987–1996, 2004.
- Sohn, B.-J., Schmetz, J., Stuhlmann, R., and Lee, J.-Y.: Dry bias in satellite-derived clear-sky water vapor and its contribution to longwave cloud radiative forcing, *J. Climate*, 19, 5570–5580, 2006.
- Stephens, G. L., Tsay, S. C., Stackhouse Jr., P. W., and Platau, P. J.: The relevance of the microphysical and radiative properties of cirrus clouds to climate and climate feedback, *J. Atmos. Sci.*, 47, 1742–1753, 1990.
- Stephens, G. L., Wood, N. B., and Gabriel, P. M.: An assessment of the parameterization of subgrid-scale cloud effects on radiative transfer. Part I: Vertical overlap, *J. Atmos. Sci.*, 61, 715–732, 2004.
- Sun, Z. and Shine, K. P.: Studies of the radiative properties of ice and mixed-phase clouds, *Q. J. Roy. Meteor. Soc.*, 120, 111–137, 1994.
- Tang, X. and Chen, B.: Cloud types associated with the Asian summer monsoons as determined from MODIS/TERRA measurements and a comparison with surface observations, *Geophys. Res. Lett.*, 33, L07814, doi:10.1029/2006GL02600, 2006.
- Taylor, J. R.: *An Introduction to Error Analysis*, University Science Books, Mill Valley, California, USA, 1982.
- Tian, L. and Curry, J. A.: Cloud overlap statistics, *J. Geophys. Res.*, 94, 9925–9935, 1989.

Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Negative cloud forcing

B. V. Thampi and R. Roca

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Warren, S. G., Hahn, C. J., and London, J.: Simultaneous occurrence of different cloud types, *J. Clim. Appl. Meteorol.*, 24, 658–667, 1985.
- Weare, B. C.: Combined satellite- and surface-based observations of clouds, *J. Climate*, 12, 897–913, 1999.
- 5 Webb, M., Senior, C., Bony, S., and Morcrette, J.-J.: Combining ERBE and ISCCP data to assess clouds in the Hadley Centre, ECMWF and LMD atmospheric climate models, *Clim. Dynam.*, 17, 905–922, 2001.
- Wentz, F. J.: A well-calibrated ocean algorithm for SSM/I, *J. Geophys. Res.*, 102, 8703–8718, 1997.
- 10 Wielicki, B. A., Cess, R. D., King, M. D., Randall, D. A., and Harrison, E. F.: Mission to planet Earth: role of clouds and radiation in climate, *B. Am. Meteorol. Soc.*, 76, 2125–2153, 1995.
- Wyser, K. and Yang, P.: Average ice crystal size and bulk shortwave single-scattering properties of cirrus clouds, *Atmos. Res.*, 49, 315–335, 1998.
- Young, D. F., Minnis, P., Doelling, D. R., Gibson, G. G., and Wong, T.: Temporal interpolation methods for the Clouds and the Earth's Radiant Energy System (CERES) Experiment, *J. Appl. Meteorol.*, 37, 572–590, 1998.
- 15 Zhang, M. H., Lin, W. Y., Klein, S. A., Bacmeister, J. T., Bony, S., Cederwall, R. T., Del Genio, A. D., Hack, J. J., Loeb, N. G., Lohmann, U., Minnis, P., Musat, I., Pincus, R., Stier, P., Suarez, M. J., Webb, M. J., Wu, J. B., Xie, S. C., and Yao, M.-S.: Comparing clouds and their seasonal variations in 10 atmospheric general circulation models with satellite measurements, *J. Geophys. Res.*, 110, D15S02, doi:10.1029/2004JD005021, 2005.
- 20 Zhang, Y., Macke, A., and Albers, F.: Effect of crystal size spectrum and crystal shape on stratiform cirrus radiative forcing, *Atmos. Res.*, 52, 59–75, 1999.
- Zhang, Y. C., Rossow, W. B., and Lacis, A. A.: Calculation of surface and top of atmosphere radiative fluxes from physical quantities based on ISCCP datasets. 1. Method and sensitivity to input data uncertainties, *J. Geophys. Res.*, 100, 1149–1165, 1995.
- 25 Zhang, Y.-C., Rossow, W. B., Lacis, A. A., Oinas, V., and Mishchenko, M. I.: Calculation of radiative fluxes from the surface to top of atmosphere based on ISCCP and other global datasets: refinements of the radiative transfer model and the input data, *J. Geophys. Res.*, 109, D19105, doi:10.1029/2003JD004457, 2004.
- 30 Zuluaga, M. D., Hoyos, C. D., and Webster, P. J.: Spatial and temporal distribution of latent heating in the south Asian monsoon region, *J. Climate*, 23, 2010–2029, doi:10.1175/2009JCLI3026.1, 2010.

Negative cloud forcing

B. V. Thampi and R. Roca

Table 1. Mean values of TOA flux and CRF (in W m^{-2}) estimated for the three negative NETCRF regimes for the summer monsoon season of 2002–2005.

	Bay of Bengal	Myanmar	Western Ghats
Clear SW	45.6	64.5	61.9
Total SW	159	173.6	156.1
Clear LW	276.7	274.9	282.7
Total LW	194.1	202.6	219.7
SWCRF	−113.4	−109.1	−94.2
LWCRF	82.6	72.3	63
NETCRF	−30.8	−36.8	−31.2

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Negative cloud forcing

B. V. Thampi and R. Roca

Table 2. Estimated uncertainties (in W m^{-2}) in the mean CERES TOA SW and LW fluxes due to various error sources for the Bay of Bengal (BOB), Myanmar (MYN) and the Western Ghats (WGS) region.

Flux	Calibration	Algorithm	Sampling uncertainty (W m^{-2})			Total uncertainty (W m^{-2})		
			BOB	MYN	WGS	BOB	MYN	WGS
TSW	1	1	1.3	1.8	3.6	1.9	2.3	3.8
TLW	1.3	1.9	0.9	2.1	1.8	2.4	3.1	2.9
CSW	0.5	1.6	2.	2.9	3.7	2.6	3.4	4.1
CLW	1	3.3	1.03	1.05	1.2	3.6	3.6	3.7

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Negative cloud forcing

B. V. Thampi and R. Roca

Table 3. Estimated uncertainty (in W m^{-2}) in the seasonal mean CERES NETCRF over the Bay of Bengal, Myanmar and the Western Ghats.

Region of Interest	Contribution of 1st term in Eq. (6)	Contribution of 2nd term in Eq. (6)	Total Uncertainty (δy) in NETCRF (W m^{-2})
Bay of Bengal	29.6	−18.1	3.4
Myanmar	39.2	−19.7	4.4
Western Ghats	52.9	−18.6	5.8

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Negative cloud forcing

B. V. Thampi and R. Roca

Table 4. Seasonal mean values of cloud micro/macro physical properties and environmental variables over the three negative NETCRF regimes for the period of June–September (2002–2005).

Region	Cloud cover (%) and cloud optical depth				Cloud particle radius (mm)		FTH	PW	Rain
	High	Upper middle	Lower middle	Lower	Ice	Water	(%)	(cm)	(mm h ⁻¹)
Bay of Bengal (10–22° N, 85–100° E)	51.6 (14.8)	25.6 (5)	4.7 (3.2)	5 (2.5)	24.8	13.9	42.2	5.7	0.42
Western Ghats (10–20° N, 72–77° E)	22 (10)	34.1 (7.2)	13.5 (7.1)	14 (5)	24.6	12.7	36.9	4.8	0.36
Myanmar (15–20° N, 92–100° E)	32.4 (12)	43.2 (8.8)	12.5 (7)	4.5 (4.8)	23.2	13.3	37.8	5.6	0.38

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Negative cloud forcing

B. V. Thampi and R. Roca

Table 5. Inter comparison of CERES TOA flux (**bold**) with that estimated from RRTM simulations (*italicized*) and ISSCP FD flux (third) for the Bay of Bengal, Myanmar and the Western Ghats regions, respectively.

TOA flux	Bay of Bengal	Myanmar	Western Ghats
Clear SW (W m ⁻²)	45.6	64.5	61.9
	<i>46.8</i>	<i>63.4</i>	<i>61.9</i>
	55.4	71	68.6
Total SW (W m ⁻²)	159	173.6	156.1
	<i>160.1</i>	<i>166</i>	<i>151.4</i>
	162.6	181.9	154.1
Clear LW (W m ⁻²)	276.7	274.9	282.7
	<i>276.9</i>	<i>275.1</i>	<i>284.7</i>
	278.7	276.2	280.5
Total LW (W m ⁻²)	194.14	202.6	219.7
	<i>195.9</i>	<i>209.6</i>	<i>225.6</i>
	206.2	215.2	231.2

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Negative cloud forcing

B. V. Thampi and R. Roca

Table 6. Inter comparison of SWCRF, LWCRF and NETCRF (in W m^{-2}) from CERES observations (**bold**) with that calculated from RRTM simulations (*italicized*) and ISCCP FD flux (third value) for the Bay of Bengal, Myanmar and the Western Ghats regions. Uncertainty associated with CERES NETCRF values are also provided (values after \pm sign).

TOA CRF	Bay of Bengal	Myanmar	Western Ghats
SWCRF	−113.4	−109.1	−94.2
	<i>−113.3</i>	<i>−102.6</i>	<i>−89.5</i>
	−107.2	−110.9	−85.5
LWCRF	82.6	72.3	63
	<i>81</i>	<i>65.5</i>	<i>59.2</i>
	72.6	61.1	49.4
NETCRF	−30.8 ± 3.4	−36.8 ± 4.4	−31.2 ± 5.8
	<i>−32.3</i>	<i>−37</i>	<i>−30.3</i>
	−34.6	−49.8	−36.1

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Negative cloud forcing

B. V. Thampi and R. Roca

Table 7. RRTM simulation of SWCRF and LWCRF values (in W m^{-2}) for the three negative NETCRF regions by changing low level cloud amount in the respective cloud vertical model.

Region		CRF (W m^{-2}) values estimated when un-obscured portions of low level clouds are	
		Present	Absent
Bay of Bengal	SWCRF	−113.4	−111.6
	LWCRF	81	80.8
Myanmar	SWCRF	−102.6	100.8
	LWCRF	66.5	66.3
Western Ghats	SWCRF	−89.5	−82.5
	LWCRF	59.2	58

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Negative cloud forcing

B. V. Thampi and R. Roca

Table 8. RRTM simulation of SWCRF (Wm^{-2}) using different ice particle size (varying SSA values) values in the cloud vertical model for the Bay of Bengal, Myanmar and the Western Ghats regions.

Ice particle radius (μm)	Bay of Bengal	Myanmar	Western Ghats
20	−116	−104.3	−90.6
25	−112.7	−102.6	−89.5
30	−109.6	−101.1	−88.5
35	−107.9	−100.2	−87.9
40	−106.2	−99.3	−87.3

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Negative cloud forcing

B. V. Thampi and R. Roca

Table 9. Seasonal mean values of TOA Flux (W m^{-2}) and CRF (W m^{-2}) over the Bay of Bengal and the West Pacific region for the period 2002–2005.

REGION	Clear Sky		Total sky		SWCRF	LWCRF	NETCRF
	SW	LW	SW	LW			
Bay of Bengal (Jun–Sep)	45.6	276.7	159	194.1	–113.4	82.6	–30.8
West Pacific (Mar–Apr)	40	285.2	107.7	220	–67.7	65.2	–2.5

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Negative cloud forcing

B. V. Thampi and R. Roca

Table 10. Seasonal mean values of cloud fraction (%) and cloud optical depth (in bracket) over the Bay of Bengal and the West Pacific region for the period 2002–2005.

REGION	Cloud cover (%) and cloud optical depth			
	High	Upper mid	Lower mid	Low
Bay of Bengal (Jun–Sep)	51.6 (14.8)	25.6 (4.9)	4.7 (3.2)	5 (2.1)
West Pacific (Mar–Apr)	38.5 (12.7)	20.1 (4.6)	4.3 (3.5)	7 (2.3)

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 1. Schematic of a typical vertical distribution of cloud fraction. Each shaded block represents the effective cloud amount for the respective cloud group estimated using the cloud fraction data and cloud overlap (maximum/random) scheme.

Negative cloud forcing

B. V. Thampi and R. Roca


[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Fig. 2. Seasonal mean variation of **(a)** SWCRF and **(b)** LWCRF over the Indian region during the summer monsoon season of 2002–2005. Color bar represents the CRF values in W m^{-2} .

Negative cloud forcing

B. V. Thampi and R. Roca




Fig. 3. Seasonal mean map of NETCRF over the Indian region during the summer monsoon season of 2002–2005. The negative NETCRF regimes (1) Bay of Bengal (2) Myanmar and (3) the Western Ghats (marked in boxes) are also shown. Color bar represents the NETCRF values in W m^{-2} .

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 4. Seasonal mean variation of **(a)** high cloud fraction (%) and **(b)** upper middle cloud fraction (%) over the Indian region during the summer monsoon season of 2002–2005. Color bar represents the mean cloud fraction in percentage.

Negative cloud forcing

B. V. Thampi and R. Roca




Fig. 5. Variation in LWCRF (left panel) and SWCRF (right panel) values with cloud top pressure for the Bay of Bengal (thick line) and Myanmar region (dashed line) respectively, estimated using RRTM.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Negative cloud forcing

B. V. Thampi and R. Roca




Fig. 6. Variation in TOA LW clear and total sky flux (left panel) and LWCRF (right panel) with Precipitable water (PW) over the Bay of Bengal region during the summer monsoon season using RRTM simulation.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 7. Contour plots of seasonal mean variation of precipitable water (mm) content over the Indian region during the (a) summer monsoon season and (b) winter season for the period 2002–2005.

Negative cloud forcing

B. V. Thampi and R. Roca




Fig. 8. Seasonal mean variation of NETCRF (W m^{-2}) over the West Pacific region during the March–April period (2002–2005). Color bar represents the NETCRF values in W m^{-2} .

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 9. Regional variation of monthly mean atmospheric water vapor (mm) from SSM/I over the Indian region (top panel) during the month of July and the West Pacific (bottom panel) during the month of April for the years 2002–2005.